

7-18-2014

Direct Comparison of Apremilast and Best Supportive Care Using a Discrete Event Simulation

Zoe Clancy, PharmD

Jefferson School of Population Health, Thomas Jefferson University, zoe.clancy@jefferson.edu

Let us know how access to this document benefits you

Follow this and additional works at: <http://jdc.jefferson.edu/msaheor>

 Part of the [Health Economics Commons](#), and the [Public Health Commons](#)

Recommended Citation

Clancy, PharmD, Zoe, "Direct Comparison of Apremilast and Best Supportive Care Using a Discrete Event Simulation" (2014). *Master of Science in Applied Health Economics and Outcomes Research*. Presentation 7.
<http://jdc.jefferson.edu/msaheor/7>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Master of Science in Applied Health Economics and Outcomes Research by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.



DIRECT COMPARISON OF APREMILAST AND BEST SUPPORTIVE CARE USING A DISCRETE EVENT SIMULATION

Zoe Clancy, PharmD

OUTLINE


- Psoriatic Arthritis Disease Brief
 - Mechanism of action
 - Differences between Psoriatic Arthritis and Rheumatoid Arthritis
 - Current treatment
- Discrete Event Simulation
 - Definition
 - Model Overview
 - Model Results



The left side of the slide features a series of vertical stripes in shades of brown, tan, and grey. Overlaid on these stripes are several orange circles of varying sizes, arranged in a cluster that tapers towards the bottom.

PSORIATIC ARTHRITIS DISEASE BRIEF

PSA IS A CHRONIC INFLAMMATORY DISEASE OF THE JOINTS AND SKIN RESULTING FROM AN UNCONTROLLED IMMUNE RESPONSE¹



Over-production of $\text{TNF-}\alpha$ as well as other cytokines, alters bone homeostasis, resulting in the joint damage seen in PsA⁴

1. Schafer. *Biochem Pharmacol.* 2012;83:1583
2. Serezani et al. *Am J Respir Cell Mol Biol.* 2008;39:127
3. Gottlieb et al. *J Am Acad Dermatol.* 2008;58:851
4. Mensah et al. *Curr Rheumatol Rep.* 2008;10(4):311

PSA DIFFERS FROM RHEUMATOID ARTHRITIS (RA) BASED ON THE PRESENCE OF PSORIATIC-ASSOCIATED CONDITIONS AND THE DISTRIBUTION AND APPEARANCE OF THE AFFECTED JOINTS

Clinical Feature	PsA	RA
Psoriatic skin lesions present	Common	No
Psoriasis-associated nail symptoms	Common	No
Distribution of affected joints	Often asymmetrical Various joints affected	Symmetrical Primarily involving hands and wrists
Appearance of the affected joint	More generalized swelling Produce a sausage-like appearance in fingers or toes	Pronounced swelling over joints (RA nodules)
Disease progression	Variable	Predictable
Rheumatoid factor status	Seronegative	Seropositive

KEY CONCEPTS

- In 75% of cases, psoriasis precedes the joint disease.
- In 15% of cases, the onset of skin disease is at the same time as onset of joint involvement.
- In 10% of cases, the joint disease precedes the psoriasis.



PSA AND QoL

- For people with psoriatic arthritis, quality of life is impacted by both the physical symptoms of the disease and the emotional burden of sometimes disfiguring skin symptoms.
- Compared to rheumatoid arthritis and ankylosing spondylitis, people with psoriatic arthritis report more psychosocial problems.
- This finding fits with data from a survey of people with psoriasis, which found that 75 percent of patients believe the skin condition had a moderate to large negative impact on their quality of life, with alteration in their activities and work.

PSA HAS A SIGNIFICANT NEGATIVE IMPACT ON HEALTH-RELATED QUALITY OF LIFE (HRQoL)

- Decreased QoL as measured by the Medical Outcomes Short-Form 36 Questionnaire (SF-36) scores in patients with PsA compared to the general population:¹
- 19% of patients with PsA claimed their disease resulted in “marked physical limitations”²
- 8.2% of patients sought assistance for home activities from friends or family³
- Both physical functioning and emotional well-being are decreased.
- In patients with PsA and psoriasis:
 - Arthritis component - greater impact on physical functioning
 - Psoriasis component - greater impact on emotional well-being
 - Skin lesions associated with poor self-image and distress from pruritus and pain.

⁸Husted JA, et al. *J Rheumatol*. 1997 Mar;24(3):511-7.

⁹Torre Alonso JC, et al. *Br J Rheumatol*. 1991 Aug;30(4):245-50.

¹⁰Kimball AB, et al. *J Drugs Dermatol*. 2007 Mar;6(3):299-306.



EPIDEMIOLOGY

Prevalence	5% - 40% of people with psoriasis
Race	Affects Caucasians more than other races
Gender	Men and women equally affected
Age of onset	40–50 years of age, can occur earlier



TREATMENT OPTIONS

Mild Disease


- NSAIDs

Moderate to Severe Disease

- Corticosteroids
- Traditional DMARDs
 - MTX
 - Sulfasalazine
 - Leflunomide
- Biologic DMARDs



National Guideline Recommendations in Patients with PsA



*No evidence supporting DMARDs ahead of TNF inhibitors (effect size: TNF inhibitors > traditional DMARDs). However, TNF inhibitors are recommended for patients who fail to respond to at least one DMARD therapy unless poor prognosis present.

Grade A=Based on evidence from meta-analysis of randomized controlled trials (RCT) or ≥ 1 RCT

Adverse Effects Limit the Benefits of Therapy with Traditional Systemic DMARDs and Biologics

■ Traditional systemic agents

- Methotrexate (MTX) has weak and conflicting evidence in the management of PsA with risks of serious toxic reactions.
- MTX is not approved by the FDA
- Leflunomide does not have FDA approval and requires monitoring for hepatic toxicity
- Sulfasalazine has limited evidence in the management of PsA with rarely occurring serious toxicities.

■ Biologics

- Mild injection-site/infusion reactions
- Black box warning of risk of serious infections and malignancies
 - Increased risk of infection
 - Overall infections, odds ratio 1.18 (95% confidence interval, 1.05-1.33)²
 - Patients with PsA are at greater risk for mortality from infection.



The Significant Burden Associated with PsA

■ **Patients with PsA:**

- Suffer from limited mobility, pain, inflammation and stiffness as well as skin lesions from psoriasis
- Have a poorer quality of life
- Are less likely to be employed and less likely to be productive
- Incur higher healthcare costs

■ **New PsA therapies are needed that demonstrate:**

- Effective Treatment in Patients with Active Psoriatic Arthritis
- Improved Safety and Better Tolerability than Traditional DMARDS and Biologics
- Patient Convenience over Injectable Biologics
- Cost savings compared to Biologics



APREMILAST

- Apremilast is a first-in-class PDE4 inhibitor
 - MOA: modulates pro-inflammatory and anti-inflammatory mediators
 - Administration: oral and does not need dose adjustments
- This drug represents a novel treatment option for patients and can represent a delay in biologic therapy¹⁴

14. Tencer T, et al. (2014) Economic evaluation of sequencing strategies in the treatment of psoriatic arthritis in the United States (abstract). Value in Health (17)3: A46.



OUTCOME MEASURES OF PsA

- ACR response criteria: 20%, 50%, 70%
(validated in RA, not PsA)
 - Tender and swollen joint count
(modified for PsA to include DIP and CMC joints: 78/76, 68/66)
 - 3/5: patient global, physician global, patient pain, HAQ, acute phase reactant (sed rate, CRP)
- Psoriatic Arthritis Response Criteria (PsARC)
 - Improvement in at least 2 of 4 criteria, including:
 - Physician global assessment (0–5)
 - Patient global assessment (0–5)
 - Tender joint score ($\geq 30\%$)
 - Swollen joint score ($\geq 30\%$)
 - Improvement in at least 1 of 2 joint scores
 - No worsening in any criteria





DISCRETE EVENT SIMULATION

All Models Are Wrong, But
Some Are Useful

-George E.P. Box



DISCRETE EVENT SIMULATION (DES)

- DES is a modeling technique that is event-based

Advantages vs Markov Models

- DES can incorporate new data as it becomes available
- Can use an individual patient's values and examine the decision from his or her point of view
- Can capture multiple events with competing risks



BASIC MODEL STRUCTURE


- Patients with active psoriatic arthritis who have failed methotrexate (MTX) therapy will be split into two groups: apremilast followed by best supportive care (BSC) and patients only receiving BSC



STEP 1: CREATE PATIENTS AND ASSIGN CHARACTERISTICS


○ Assign Baseline Utilities:

- Age
- Gender (45% male)
- Life expectancy
- Mortality




STEP 2: PATIENTS ENTER EITHER APR OR BSC TREATMENT ARM

- If ‘Is Patient starting a Trial?’ is TRUE, then patients enter APR arm
- ‘Assign Time to Event TP’ sets the next event to death and logs the time at the beginning of the time-to-event period
- Time advances in “Wait Next Event TP”




STEP 2: PATIENTS ENTER EITHER APR OR BSC TREATMENT ARM

- Patients move to 'Assign QALYs and Costs TP' where QALYs and Costs are calculated
- The VBA module is used to calculate Other Healthcare Costs
 - The VBA module computes the patient's age each month and tallies the costs over the course of the period




STEP 2: PATIENTS ENTER EITHER APR OR BSC TREATMENT ARM

- After costs and QALYs have been assigned, 'Death in TP?' checks to see if the time of death event was prior to the end of the Trial Period
- If so, patient is disposed of in the model, otherwise patient continues to BSC




STEP 3: DECIDE IF TREATMENT WAS EFFECTIVE (OR NOT)

- Patients enter a decision module ('DECIDE outcome of Trial Period') which decides whether the patients achieved a PsARC score (effective treatment) or not
- If treatment effective, patients are assigned to a PASI group to allocate future costs and QALYs
- Patients who are not successfully treated move to the BSC arm




STEP 3: DECIDE IF TREATMENT WAS EFFECTIVE (OR NOT)

- 'Assign Time to Event PASI' module assigns a length of time until patients move to BSC
- Similar to the Trial Period arm, patient is advanced in time through the 'Wait Next Event PASI' module to the sooner of either Death or BSC or model end
- Costs and QALYs are assigned as in Trial Period arm



STEP 4: BSC, DEATH, OR MODEL END

- Patients enter BSC arm either at beginning of the model run or through discontinuation of treatment
- Similar to Trial Period and Apremilast Arm, with patients disposed of at the end
- The Excel read/write modules are also shown



MODEL ASSUMPTIONS

- Patients who enter the BSC arm do not go back to apremilast therapy
- There are no changes to BSC or treatment paradigms of PsA in clinical practice over the time horizon of the model (5 years)
- The population to which the model is applied to follows the age and life expectancy of that in the model
- HAQ scores return to baseline after discontinuation of treatment
- No monitoring or lab costs for apremilast



MODEL LIMITATIONS

- Data was sourced from clinical trials and not real world
- PASI is used as the trial period endpoint, but is not the clinical trial endpoint for efficacy
- Indirect costs of treatment are not accounted for in the model



MODEL RESULTS

		Control		Apremilast		
Rep		Costs	QALYs	Costs	QALYs	
1.00		\$ 30,558.54	19.70	\$ 288,081.27	34.00	
2.00		\$ 107,698.74	53.81	\$ 162,649.05	54.36	
3.00		\$ 66,412.11	36.19	\$ 302,328.11	48.89	
4.00		\$ 139,131.43	68.18	\$ 651,137.37	74.52	
5.00		\$ 32,584.15	17.74	\$ 128,977.00	20.67	
6.00		\$ 86,104.37	45.88	\$ 188,795.84	47.13	
7.00		\$ 75,148.61	38.87	\$ 242,798.90	40.83	
8.00		\$ 94,376.35	50.68	\$ 198,235.75	53.69	
9.00		\$ 99,138.22	53.90	\$ 224,178.43	60.20	
10.00		\$ 96,838.01	57.71	\$ 226,460.36	59.30	
11.00		\$ 75,393.11	41.54	\$ 210,705.91	48.16	
12.00		\$ 106,148.80	55.74	\$ 504,847.20	60.29	
13.00		\$ 103,474.64	50.81	\$ 123,192.79	51.13	
14.00		\$ 73,509.76	37.78	\$ 142,757.01	38.63	
15.00		\$ 113,050.81	64.10	\$ 246,031.07	65.72	
	Total	\$ 1,299,567.67	692.65	\$ 3,841,176.07	757.51	
	Avg/Patient	\$ 17,327.57	9.24	\$ 51,215.68	10.10	
			AVG	\$ 33,888.11	0.86	

Comparison of DES to Markov Model

- Model cost results are within 20%

	QALY/ patient	Cost/ patient
Markov Model	0.29	\$41,338
DES	0.86	\$33,888



CONCLUSIONS

- DES models are more adaptable, compared to Markov models
 - Once data becomes available, for example QOL instrument data, the DES is easily updated
- DES and Markov models share limitations, specifically the availability and quality of data
 - Markov models have less data requirements
- A comparison of two models with the same data shows differences that can be attributed to
 - time to event that was used to calculate drop off to BSC
 - distributions used for age and life expectancy



REFERENCES

1. Schafer. *Biochem Pharmacol*. 2012;83:1583
2. Serezani et al. *Am J Respir Cell Mol Biol*. 2008;39:127
3. Gottlieb et al. *J Am Acad Dermatol*. 2008;58:851
4. Mensah et al. *Curr Rheumatol Rep*. 2008;10(4):311
5. Husted JA, et al. *Arthritis Rheum*. Apr 2001;45(2):151-158.
6. Salaffi F et al. *Health Qual Life Outcomes*. 2009;7:25.
7. Bhosle MJ, et al. *Health Qual Life Outcomes*. 2006;4:35.
8. Husted JA et al. *J Rheumatol*. 1997 Mar;24(3):511-7.
9. Torre Alonso JC et al. *Br J Rheumatol*. 1991 Aug;30(4):245-50.
10. Kimball AB et al. *J Drugs Dermatol*. 2007 Mar;6(3):299-306
11. Gladman DD. *Rheum Dis Clin North Am* 1992;18:247–56
12. Ritchlin CT et al. *Ann Rheum Dis*. 2009 Sept;68(9): 1387-94.
13. Enbrel Prescribing Information,
http://pi.amgen.com/united_states/enbrel/derm/enbrel_pi.pdf
14. Tencer T et al. *Value in Health* 2014 (17)3: A46.

